



### Nuclear Stress Test Order

**Patient:** \_\_\_\_\_ **Date of Study:** \_\_\_\_\_

**ID#:** \_\_\_\_\_ **DOB:** \_\_\_\_\_ **Weight:** \_\_\_\_\_

**Physician:** \_\_\_\_\_ **Test:** Exercise Lexiscan Dobutamine  
MUGA (\* circle test ordered)

**Test Indication:** \_\_\_\_\_

**Females** (age: 18 to 55): Y / N      Hysterectomy: Y / N      HCG ordered \*: Y / N  
Mastectomy: Y / N      Breast prosthesis or implants: Y / N  
Breastfeeding: please notify physician before having procedure.  
**\*HCG results needed 24 hrs prior to test** on females < 55 years old without  
previous hysterectomy or uterine ablation.

**Additional clinical information:** \_\_\_\_\_  
Pre-op clearance: Y / N      Surgery: \_\_\_\_\_  
Date surgery scheduled: \_\_\_\_\_ Surgeon: \_\_\_\_\_  
Pacemaker/AICD: Y / N      LBBB on EKG: Y / N  
Abd. Aortic Aneurysm: Y / N (cm: \_\_\_\_\_)      Aortic stenosis: Y / N  
Coronary disease: Y / N      Prior CABG: Y / N  
Coronary stent: Y / N

**Medications:** (please list)  
**Beta blockers:** \_\_\_\_\_  
(\* **Beta blocker therapy** suggested to be held 24 hours prior if stress test being  
done for diagnosis.)  
Theophylline, or theophylline-like meds: \_\_\_\_\_  
Aggrenox, or meds containing Persantine/Dipyridamole: \_\_\_\_\_  
(\*\* **Theophylline** interferes with Lexiscan vasodilation-- med needs to be held  
12 hrs. prior to procedure. **Persantine/Dipyridamole** may increase the activity of  
Lexiscan--med needs to be held 48 hrs prior to procedure)  
Oxygen per N/C?: \_\_\_\_\_

**Received Nuclear Stress Test Patient Instructions?** Y / N

Checklist completed by: \_\_\_\_\_ for \_\_\_\_\_, MD/DO

(Date) \_\_\_\_\_